

ON THE DISTRIBUTION OF COEFFICIENTS OF RESIDUE POLYNOMIALS

LÁSZLÓ MAJOR

ABSTRACT. For any nonconstant polynomial with positive coefficients and positive integer d if $r(p^n)$ denotes the least degree residue of p^n modulo $x^d - 1$, then the coefficients of $r(p^n)$ are asymptotically equal as n tends to infinity.

When Euler (1765) was investigating the properties of the trinomial coefficients (see Andrews [1]), he obtained an unimodal distribution of the coefficients by expanding the n^{th} power of the polynomial $1 + x + x^2$. In the analogous case of the polynomial $1 + x$ the corresponding unimodal distribution is the n^{th} row of Pascal's triangle. In general the distribution of the coefficients of the n^{th} power of the polynomial $p = p_0 + p_1x + \cdots + p_mx^m$ with nonnegative real coefficients is not necessarily unimodal, but a sufficient condition is given for unimodality by Boros and Moll [2]. We propose to examine the coefficient distribution of the powers of p in the polynomial residue class ring modulo $x^d - 1$.

Let $2 \leq d$ be a fixed positive integer. We consider the polynomial ring $\mathbb{R}[x]$ and the quotient ring $\mathbb{R}[x]/x^d - 1$. Let p denote both a polynomial of $\mathbb{R}[x]$ with nonnegative coefficients and the residue class of $\mathbb{R}[x]/x^d - 1$ represented by p . The proper meaning will be determined by the context. We use the most common representative system formed by all polynomials of degree less than d . Let $r(p)$ denote that unique polynomial of degree less than d (denoted $\deg(r(p)) < d$) for which the polynomial $p \in \mathbb{R}[x]$ is congruent to $r(p)$ modulo $x^d - 1$. In consequence the n^{th} power of $p \in \mathbb{R}[x]/x^d - 1$ will mean the residue class of $r(p^n)$. A polynomial with positive (nonnegative) coefficients is said to be *positive (nonnegative)* polynomial.

The main result is Theorem 1. The lemmas and corollaries provide background and an extension.

Lemma 1. *Let $p \in \mathbb{R}[x]$ be a nonnegative, nonzero polynomial. For any $q \in \mathbb{R}[x]$ and $i \in \mathbb{N}$ let q_i denote the coefficient of x^i in q . If there exist $k, l \in \{0, \dots, d-1\}$ such that $r(p)_k > 0$, $r(p)_l > 0$ and $\gcd(d, k-l) = 1$, then $r(p^{d-1})$ is a positive polynomial.*

Proof. It can be assumed that $k < l$. Let us denote the difference $l - k$ by h . It is easy to see that if $r((x^k + x^l)^{d-1})$ is positive then also $r(p^{d-1})$ is positive. In addition, $r((x^k + x^l)^{d-1}) = r(x^{k(d-1)}(1 + x^h)^{d-1})$ is positive if $r((1 + x^h)^{d-1})$ is positive, so let us concentrate on the polynomial $r((1 + x^h)^{d-1})$. We expand the

Date: May 19, 2008.

Key words and phrases. coefficient distribution, polynomial residue class ring, positive polynomial, circulant matrix.

expression $(1 + x^h)^{d-1}$ in the polynomial ring $\mathbb{R}[x]$ by using the Binomial Theorem:

$$(0.1) \quad (1 + x^h)^{d-1} = \sum_{i=0}^{d-1} \binom{d-1}{i} (x^{hi}).$$

Let us assume that the following congruence holds for some $i, j \in \{0, \dots, d-1\}$:

$$(0.2) \quad \binom{d-1}{i} (x^{hi}) \equiv \binom{d-1}{j} (x^{hj}) \pmod{x^d - 1}.$$

From the equation (0.2) follows that $hi - hj$ is divisible by d . But it was assumed that $\gcd(h, d) = 1$, consequently $i = j$. Therefore $r(p^{d-1})$ has exactly d different nonzero coefficients, in other words it is positive. $\square$

Corollary 1. *If $n \geq d-1$ is an integer and p satisfies the conditions of the Lemma 1, then $r(p^n)$ is a positive polynomial.*

In order to be able to deal more efficiently with powers of polynomials we need the concept and some useful properties of *circulant matrices*. For a classical reference see e.g. Davis [3]. Let $v = (v_0, v_1, \dots, v_{d-1})$ be a row vector in $\mathbb{R}^d$. The permutation $\rho : \mathbb{R}^d \rightarrow \mathbb{R}^d$ given by

$$\rho(v_0, v_1, \dots, v_{d-1}) = (v_{d-1}, v_0, \dots, v_{d-2})$$

is called *cyclic permutation*. The *circulant matrix* associated to the vector v is the $d \times d$ matrix whose i^{th} row is $\rho^{i-1}(v)$, $i = 1, \dots, d$ and it is denoted by

$$C = \text{circ}(v_0, v_1, \dots, v_{d-1}) = \text{circ}(v).$$

The product of two circulant matrices is circulant, therefore any positive integer power of a circulant matrix is circulant. We shall use the connection between powers of circulant matrices and the powers of residue polynomials. If $a = (a_0, a_1, \dots, a_{d-1})$ is the coefficient vector of some polynomial $r(p) \in \mathbb{R}[x]$, then the first row of $(\text{circ}(a))^n$ is the coefficient vector of $r(p^n)$. Lally and Fitzpatrick [4] give a general overview in this subject.

A matrix is called *positive (nonnegative)* if all its entries are positive (nonnegative). A $d \times d$ nonnegative matrix M is said to be *doubly stochastic* if the sum of the entries in each row and in each column equals 1. The product of doubly stochastic matrices is doubly stochastic. The following lemma follows from a general result on eigenvectors of positive matrices, Theorem 8.2.8 of Horn and Johnson [5]. Here we provide a direct proof. A matrix whose entries are all ones will be denoted by U .

Lemma 2. *If M is a $d \times d$ positive doubly stochastic matrix, then*

$$(0.3) \quad \lim_{n \rightarrow \infty} M^n = \frac{1}{d} U.$$

Proof. Let us denote the doubly stochastic matrix $\frac{1}{d}U$ by J . If $M = J$, then the statement is trivial. Otherwise the matrix M can be given in the following form:

$$(0.4) \quad M = \lambda J + (1 - \lambda) M_0,$$

where $0 < \lambda < 1$ and M_0 is a doubly stochastic matrix. Let us write the n^{th} power of M by using the Binomial Theorem and the fact that $JD = DJ = J$ for all

doubly stochastic matrix D ,

$$\begin{aligned} M^n &= \sum_{i=0}^n \binom{n}{i} (\lambda J)^{n-i} ((1-\lambda)M_0)^i \\ &= (\lambda J)^n + J \sum_{i=1}^n \binom{n}{i} \lambda^{n-i} (1-\lambda)^i = (\lambda)^n J + J. \end{aligned}$$

From the above expression follows our statement, because $0 < \lambda < 1$ and consequently $\lim_{n \rightarrow \infty} (\lambda)^n = 0$. $\square$

In order that the Lemma 2 could be applied for the powers of polynomials, we shall restrict our attention to polynomials p whose coefficients sum to 1 ($p(1) = 1$). We may do this without loss of generality, because all polynomials can be written in the form $p = p(1) \cdot p'$, where p' has the mentioned property. It can also be said that the coefficients of p' form a *stochastic vector*. Clearly the sum of the coefficients of $(p')^n$ is also 1 and the sum of the coefficients of p^n is equal to $p(1)^n$. By using Lemma 1, Lemma 2 and the connection between powers of circulant matrices and the powers of residue polynomials we obtain the following theorem:

Theorem 1. *For any polynomial q and nonnegative integer i let q_i denote the coefficient of x^i in q . Let p be a nonconstant polynomial with positive real coefficients summing to 1 and $d \geq 2$ a fixed integer. Then for all $0 \leq j \leq d-1$*

$$(0.5) \quad \lim_{n \rightarrow \infty} \sum_{k \equiv j \pmod{d}} (p^n)_k = \frac{1}{d}$$

The restriction, that p is positive, is quite strong. Theorem 1 can be extended to larger classes of polynomials. For instance the following corollary applies Lemma 1 so as to provide such an extension.

Corollary 2. *If the nonnegative polynomial p satisfies the conditions of Lemma 1 and its coefficients sum to 1, then the convergence (0.5) holds for p or equivalently for all $0 \leq j \leq d-1$*

$$\lim_{n \rightarrow \infty} r(p^n)_j = \frac{1}{d}.$$

Proof. Let p be a polynomial satisfying the conditions of Lemma 1 and let $p(1) = 1$. Let us denote the coefficient vector of $r(p)$ by c and the circulant $\text{circ}(c)$ by C . By Corollary 1 if $m \geq d-1$ then C^m is a positive matrix. We use J to denote the $d \times d$ matrix in which every element is $\frac{1}{d}$. By Lemma 2, $\lim_{n \rightarrow \infty} C^{(d-1)n} = J$. We may also express this fact by using the max norm of matrices:

$$\lim_{n \rightarrow \infty} \|C^{(d-1)n} - J\| = 0.$$

We show that for all $0 \leq i \leq d-1$, $\lim_{n \rightarrow \infty} C^{(d-1)n+i} = J$. We recall that $JC^i = J$, so we have

$$(0.6) \quad \|C^{(d-1)n}C^i - J\| = \|C^{(d-1)n}C^i - JC^i\| \leq \|C^{(d-1)n} - J\| \cdot \|C^i\|.$$

Because $\|C^i\|$ is a constant, we obtain that $\lim_{n \rightarrow \infty} C^{(d-1)n+i} = J$ for all $0 \leq i \leq d-1$, consequently $C^m \rightarrow J$ as m tends to infinity. $\square$

Example Theorem 1 applies to the polynomial $p = 1 + x$ and $d = 3$. The n^{th} row of Pascal's triangle displays the coefficients of p^n . In this case Theorem 1 says that the sum of every third element of the n^{th} row of Pascal's triangle is asymptotically equal to $\frac{1}{3} \cdot 2^n$.

REFERENCES

- [1] G.E. ANDREWS *Euler's "Exemplum Memorabile Inductionis Fallacis" and q -Trinomial Coefficients*, J. American Math. Soc., 3 (1990), 653-669.
- [2] G. BOROS, V. MOLL, : *A Criterion for Unimodality*, The Electronic Journal of Combinatorics, **6**, (1999), 1-6.
- [3] P.J. DAVIS: *Circulant Matrices*, John Wiley and Sons, 1979.
- [4] K. LALLY, P. FITZPATRICK: *Algebraic Structure of Quasicyclic Codes*, Discrete Applied Mathematics, **111**, (2001), 157-175.
- [5] R.A. HORN, C.R. JOHNSON: *Matrix Analysis*, Cambridge University Press, 1991.

LÁSZLÓ MAJOR
INSTITUTE OF MATHEMATICS
TAMPERE UNIVERSITY OF TECHNOLOGY
FINLAND
E-mail address: laszlo.major@tut.fi